

# Full left ventricular coverage is essential for the accurate quantification of the area-at-risk by T1 and T2 mapping

Bulluck H., Bryant J., Lim M., Tan X., Ramlall M., Francis R., Kotecha T., Cabrera-Fuentes H., Knight D., Fontana M., Moon J., Hausenloy D.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© 2017 The Author(s). T2-weighted cardiovascular magnetic resonance (CMR) using a 3-slice approach has been shown to accurately quantify the edema-based area-at-risk (AAR) in ST-segment elevation myocardial infarction (STEMI). We aimed to compare the performance of a 3-slice approach to full left ventricular (LV) coverage for the AAR by T1 and T2 mapping and MI size. Forty-eight STEMI patients were prospectively recruited and underwent a CMR at  $4 \pm 2$  days. There was no difference between the AAR full LV and AAR 3-slices by T1 ( $P = 0.054$ ) and T2-mapping ( $P = 0.092$ ), with good correlations but small biases and wide limits of agreements (T1-mapping:  $N = 30$ ,  $R^2 = 0.85$ , bias =  $1.7 \pm 9.4\%$  LV; T2-mapping:  $N = 48$ ,  $R^2 = 0.75$ , bias =  $1.7 \pm 12.9\%$  LV). There was also no significant difference between MI size 3-slices and MI size full LV ( $P = 0.93$ ) with an excellent correlation between the two ( $R^2 = 0.92$ ) but a small bias of  $0.5\%$  and a wide limit of agreement of  $\pm 7.7\%$ . Although MSI was similar between the 2 approaches, MSI 3-slices performed poorly when MSI was  $< 0.50$ . Furthermore, using AAR 3-slices and MI size full LV resulted in 'negative' MSI in 7/48 patients. Full LV coverage T1 and T2 mapping are more accurate than a 3-slice approach for delineating the AAR, especially in those with MSI  $< 0.50$  and we would advocate full LV coverage in future studies.

<http://dx.doi.org/10.1038/s41598-017-05127-0>

---

## References

- [1] Friedrich, M. G. et al. The salvaged area at risk in reperfused acute myocardial infarction as visualized by cardiovascular magnetic resonance. *Journal of the American College of Cardiology* 51, 1581-1587 (2008).
- [2] Botker, H. E., Kaltoft, A. K., Pedersen, S. F. & Kim, W. Y. Measuring myocardial salvage. *Cardiovascular research* 94, 266-275 (2012).
- [3] Engblom, H. et al. Sample Size in Clinical Cardioprotection Trials Using Myocardial Salvage Index, Infarct Size, or Biochemical Markers as Endpoint. *Journal of the American Heart Association* 5, Advance online publication (2016).
- [4] Schulz-Menger, J. et al. Standardized image interpretation and post processing in cardiovascular magnetic resonance: Society for Cardiovascular Magnetic Resonance (SCMR) board of trustees task force on standardized post processing. *Journal of cardiovascular magnetic resonance: official journal of the Society for Cardiovascular Magnetic Resonance* 15, 35 (2013).
- [5] Bulluck, H. et al. T1 mapping and T2 mapping at 3T for quantifying the area-at-risk in reperfused STEMI patients. *Journal of cardiovascular magnetic resonance: official journal of the Society for Cardiovascular Magnetic Resonance* 17, 73 (2015).

- [6] Dall'Armellina, E. et al. Cardiovascular magnetic resonance by non contrast T1-mapping allows assessment of severity of injury in acute myocardial infarction. *Journal of cardiovascular magnetic resonance: official journal of the Society for Cardiovascular Magnetic Resonance* 14, 15 (2012).
- [7] Berry, C. et al. Magnetic Resonance Imaging Delineates the Ischemic Area at Risk and Myocardial Salvage in Patients With Acute Myocardial Infarction. *Circulation-Cardiovascular Imaging* 3, 527-535 (2010).
- [8] Kim, H. W. et al. Relationship of T2-Weighted MRI Myocardial Hyperintensity and the Ischemic Area-At-Risk. *Circulation research* 117, 254-265 (2015).
- [9] Giri, S. et al. T2 quantification for improved detection of myocardial edema. *Journal of cardiovascular magnetic resonance: official journal of the Society for Cardiovascular Magnetic Resonance* 11, 56 (2009).
- [10] Verhaert, D. et al. Direct T2 quantification of myocardial edema in acute ischemic injury. *JACC. Cardiovascular imaging* 4, 269-278 (2011).
- [11] Ugander, M. et al. Myocardial edema as detected by pre-contrast T1 and T2 CMR delineates area at risk associated with acute myocardial infarction. *JACC. Cardiovascular imaging* 5, 596-603 (2012).
- [12] Hamshere, S. et al. Cardiovascular magnetic resonance imaging of myocardial oedema following acute myocardial infarction: Is whole heart coverage necessary? *Journal of cardiovascular magnetic resonance: official journal of the Society for Cardiovascular Magnetic Resonance* 18, 7 (2016).
- [13] Arai, A. E. Using magnetic resonance imaging to characterize recent myocardial injury: utility in acute coronary syndrome and other clinical scenarios. *Circulation* 118, 795-796 (2008).
- [14] Pennell, D. Myocardial salvage: retrospection, resolution, and radio waves. *Circulation* 113, 1821-1823 (2006).
- [15] Langhans, B. et al. Reproducibility of area at risk assessment in acute myocardial infarction by T1-and T2-mapping sequences in cardiac magnetic resonance imaging in comparison to Tc99m-sestamibi SPECT. *The international journal of cardiovascular imaging* 30, 1357-1363 (2014).
- [16] Bulluck, H. et al. Automated Extracellular Volume Fraction Mapping Provides Insights Into the Pathophysiology of Left Ventricular Remodeling Post-Reperfused ST-Elevation Myocardial Infarction. *Journal of the American Heart Association* 5 (2016).
- [17] Bulluck, H. et al. Residual Myocardial Iron Following Intramyocardial Hemorrhage During the Convalescent Phase of Reperfused ST-Segment-Elevation Myocardial Infarction and Adverse Left Ventricular Remodeling. *Circulation: Cardiovascular Imaging* 9, e004940 (2016).
- [18] Bulluck, H. et al. Impact of microvascular obstruction on semiautomated techniques for quantifying acute and chronic myocardial infarction by cardiovascular magnetic resonance. *Open Heart* 3, e000535 (2016).
- [19] Bulluck, H. et al. Defining left ventricular remodeling following acute ST-segment elevation myocardial infarction using cardiovascular magnetic resonance. *Journal of cardiovascular magnetic resonance: official journal of the Society for Cardiovascular Magnetic Resonance* 19, 26 (2017).
- [20] Bulluck, H. et al. Diagnostic Performance of T1 and T2 Mapping to Detect Intramyocardial Hemorrhage in Reperfused ST-Segment Elevation Myocardial Infarction (STEMI) Patients. *Journal of magnetic resonance imaging: JMRI* (2017).
- [21] Steg, P. G. et al. ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *European heart journal* 33, 2569-2619 (2012).
- [22] Kellman, P. & Hansen, M. S. T1-mapping in the heart: accuracy and precision. *Journal of cardiovascular magnetic resonance: official journal of the Society for Cardiovascular Magnetic Resonance* 16, 2 (2014).
- [23] Xue, H. et al. Motion correction for myocardial T1 mapping using image registration with synthetic image estimation. *Magnetic resonance in medicine: official journal of the Society of Magnetic Resonance in Medicine/Society of Magnetic Resonance in Medicine* 67, 1644-1655 (2012).
- [24] Ledesma-Carbayo, M. J., Kellman, P., Hsu, L. Y., Arai, A. E. & McVeigh, E. R. Motion corrected free-breathing delayed enhancement imaging of myocardial infarction using nonrigid registration. *Journal of magnetic resonance imaging: JMRI* 26, 184-190 (2007).
- [25] Kellman, P. & Arai, A. E. Cardiac imaging techniques for physicians: late enhancement. *Journal of magnetic resonance imaging: JMRI* 36, 529-542 (2012).